EDINBURGH 97/3
ITP-SB-97-24

arXiv:hep-ph/9705234v1 5 May 1997

Resummation for QCD Hard Scattering

Nikolaos Kidonakis

Department of Physics and Astronomy
University of Edinburgh
Edinburgh EH9 3JZ, Scotland

George Sterman

Institute for Theoretical Physics
State University of New York at Stony Brook
Stony Brook, NY 11794-3840, USA

May 1997

Abstract

We resum distributions that are singular at partonic threshold (the elastic limit) in heavy quark production, in terms of logarithmic behavior in moment space. The method may be applied to a variety of cross sections sensitive to the edge of phase space, including transverse momentum distributions. Beyond leading logarithm, dependence on the moment variable is controlled by a matrix renormalization group equation, reflecting the evolution of composite operators that represent the color structure of the underlying hard scattering. At next-to-leading logarithmic accuracy, these evolution equations may be diagonalized, and moment dependence in the cross section is a sum of exponentials. Beyond next-to-leading logarithm, resummation involves matrix-ordering. We give a detailed analysis for the case of heavy quark production by light quark annihilation and gluon fusion.

1 Introduction

The factorization of cross sections into long- and short-distance functions gives QCD predictive power for a wide variety of inclusive, hard-scattering cross sections in hadronic collisions. There has been great progress in recent years in the calculation of perturbative corrections at next-to-leading order (NLO) for such processes, including Drell-Yan [1], direct photon [2], heavy quark [3] and jet production [4]. Selected exact results at two or even more loops are also available, especially for e^+e^- annihilation [5] and deeply inelastic scattering [6]. The exploration of quantum chromodynamics, electroweak theory, and searches for extensions of the standard model, however, require information on yet higher orders. We need to develop methods to decide when higher-order perturbative terms are negligible and, when they are not, to estimate their importance. This motivates the systematic study of pertubation theory at arbitrary order. One of the elements in this program is the resummation of singular distributions in factorized inclusive cross sections. This resummation is most conveniently carried out in moment space. Closely related resummations organize large logarithmic corrections near zero transverse momentum for heavy systems such as Drell-Yan or heavy quark pairs.

In this paper we shall give details of this resummation in hard-scattering cross sections induced by QCD itself, particularly heavy quark production, extending results for electroweak-induced hard scattering cross sections, such as the Drell-Yan production of vector bosons [7, 8, 9]. Our starting point will be the factorization properties of these cross sections near threshold, and we shall make strong use of the close relation between factorization and resummation [10].

The resummations that we consider apply to processes of the form

$$h_1(p_1) + h_2(p_2) \to T(q, y, \chi) + X \,, \tag{1.1}$$

where $T(q, y, \chi)$ represents a system in the hadronic final state, produced with large invariant mass $\sqrt{q^2} = Q$ and (collective) rapidity y,

$$y = \frac{1}{2} \ln \left(\frac{q \cdot p_2}{q \cdot p_1} \right) \,. \tag{1.2}$$

Examples include the production of heavy quark pairs and high-p_T jets. The additional variable χ represents the internal structure of the final state, for

example the angle θ between the quark, or jet, direction in the final state and the beam axis. Such cross sections may be written in factorized form as

$$\begin{aligned}\frac{d\sigma_{h_1\,h_2}}{dQ^2\,d\chi\,dy} &= \sum_{ab}\int dx_a dx_b\,\phi_{a/h_1}(x_a,\mu^2)\phi_{b/h_2}(x_b,\mu^2) \\ &\quad\times H_{ab}\left(\frac{Q^2}{x_a x_b S}, y-y_p, \chi, \frac{Q}{\mu}, Q, \alpha_s(\mu^2)\right), \end{aligned} \tag{1.3}$$

where S is the center-of-mass (c.m.) energy squared of the hadrons, and where the ϕ's are parton distributions defined in an appropriate scheme, and evaluated at factorization scale μ. The function H_{ab} is the hard-scattering partonic cross section (the "hard part") that describes the production of the final-state system T in the collision of partons a and b, which carry momentum fractions x_a and x_b of hadrons h_1 and h_2, respectively. Rapidity-dependence in H_{ab} is through the combination $y-y_p$, where y_p is the partonic rapidity in the hadronic c.m. frame,

$$y_p = \frac{1}{2}\ln\left(\frac{x_a}{x_b}\right). \tag{1.4}$$

The hard parts H_{ab} in Eq. (1.3) are calculable in perturbation theory. In general, they are not smooth functions of their arguments, but combinations of functions and singular distributions. The reason for this somewhat complicated structure is that they summarize cancellations between cross sections with gluon emission and with virtual gluon corrections. This cancellation (in QCD as in QED) is only manifest at the level of sufficiently inclusive cross sections. For factorized cross sections such as Eq. (1.3), this requirement manifests itself in the occurrence of ("plus") distributions, which are singular for $z=1$, where

$$z = \frac{Q^2}{s}, \tag{1.5}$$

with $s = x_a x_b S$ the invariant mass squared of the partons that initiate the hard scattering. We shall refer to $z=1$ as "partonic threshold" [1], or sometimes as the "elastic limit".

[1] We emphasize that by partonic threshold, we refer to the c.m. total energy of the incoming partons for a fixed final state; heavy quarks, for example, are not necessarily produced at rest.

Singular distributions are most easily organized in terms of moments with respect to $\tau = Q^2/S$. As we review below, for the Drell-Yan and related cross sections logarithmic dependence on the moment variable exponentiates [7]-[11]. This moment must be inverted to derive the physical cross section. There are a number of proposals on the best way to carry out this transformation, or otherwise use the resummation, which have been put forward in the context of Drell-Yan [12, 13, 14] and heavy quark production [15, 16, 17, 18, 19, 20]. Our analysis here will not deal directly with this question; our results will remain in moment space. In this paper, we shall be concerned with the extension of resummation to cross sections that are initiated by QCD itself. Our explicit applications will be to resummation at partonic threshold for heavy quark production, but the method is much more general. A straightforward application of the method, which involves only the replacement of the moments by Fourier transformation to impact parameter space, is to the overall transverse momentum of the heavy quark pair, again extending results for Drell-Yan pairs [21], and hadrons in e^+e^- annihilation [22].

It has been recognized for some time that leading logarithms in moment (or impact parameter) space exponentiate for heavy quark production [15, 16, 17] in the same manner as for Drell-Yan, reflecting the factorizability of leading singular distributions from the hard subprocess [7]. Beyond the level of leading logarithms, however, singular distributions in QCD-induced cross sections depend coherently on color exchange in the underlying hard scattering. The interplay between the color structures of short and long distance interactions will be a unifying thread of the discussion that follows.

The next section starts with a brief review of the factorization of the inclusive Drell Yan cross sections near partonic threshold. We shall describe the resummation of the singular distributions in these cross sections starting from their factorization properties. We also observe the field-theoretic content of the functions into which the cross section factorizes, relating them to matrix elements of fields and of Wilson lines [23, 24, 25, 26, 27]. We then extend the factorization, and hence resummation, formalism to QCD-induced inclusive cross sections. The possibility for, and the broad outlines of, such an extension has recently been discussed [10]; here, the practical formalism, and its field theoretic basis, are discussed in some detail, expanding on a summary given in Ref. [28] (see also [29]). To organize sensitivity to color exchange at short distances beyond leading order in $\ln(1-z)$ (or in loga-

rithms of the corresponding moment variable), we derive a matrix evolution equation that controls all logarithmic enhancements at threshold. At next-to-leading logarithm (NLL) in the exponent, the matrix equation may be diagonalized in a manner reminiscent of leading order evolution in deeply inelastic scattering. At NLL, the cross section is a sum of exponentials, each representing an eigenvector of the anomalous dimension matrix. Beyond this level of accuracy, however, the resummation can only be carried out in terms of ordered exponentials. Resummation at NLL requires the computation of the anomalous dimension matrix for each partonic subprocess, such as light quark annihilation into heavy quarks or gluon fusion into heavy quarks.

The third section describes the calculation of the anomalous dimension matrix for light quark annihilation into heavy quarks, supplementing the brief description of Ref. [28]. Section 4 extends these results to the very important case of gluon fusion. The paper concludes with a summary and the outlook for further progress.

2 Factorization and Resummation

2.1 Resummation for Color Singlet Hard Scattering

The resummation of singular distributions for the Drell-Yan, and related electroweak-induced hard scattering processes has been understood for some time [7, 8, 9]. In this subsection, we review the basic results, and elaborate somewhat on the operators in QCD that play a role in the underlying factorization. For our purposes, it will be useful to generalize the standard results slightly, to the production of an arbitrary color-singlet system, denoted B below, of invariant mass squared Q^2. We assume in particular that B may be produced by gluon fusion as well as quark annihilation. An important example is Higgs production [30]. For the purposes of the following discussion, however, we shall refer to this class of production processes as Drell-Yan cross sections.

Resummation is most readily formulated in terms of moments of the inclusive Drell-Yan cross section $d\sigma/dQ^2$, with respect to the variable $\tau = Q^2/S$,

$$\int_0^1 d\tau \tau^{N-1} \frac{1}{\sigma_0} \frac{d\sigma_{h_1 h_2 \to B}}{dQ^2}(\tau, Q^2) \;\; = \;\; \sum_{i=q,\bar{q},g} e_f^2 \; \tilde{\phi}_{i/h_1}(N, \mu^2) \, \tilde{\phi}_{\bar{i}/h_2}(N, \mu^2)$$

$$\times\tilde{\omega}_{i\bar{i}\to B}\left(N, Q/\mu, \alpha_s(\mu^2)\right) + \mathcal{O}(1/N)\,. \tag{2.1}$$

Here, e_f is the fractional quark charge, σ_0 ($= 4\pi\alpha^2/(9Q^2S)$ for the standard Drell-Yan cross section) is a Born cross section, and the moment functions $\tilde{\phi}$ are

$$\tilde{\phi}(N,\mu^2) = \int_0^1 dx\; x^{N-1}\phi(x,\mu^2)\,, \tag{2.2}$$

and similarly for the hard-scattering function $\omega_{i\bar{i}\to B}(z, Q/\mu, \alpha_s(\mu^2))$ in terms of the variable z. In Eq. (2.1) we have neglected hard-scattering functions associated with the gluon-quark combination, with two quarks or antiquarks, and with quarks and antiquarks of different flavors, since these do not give rise to singular distributions at $z = 1$. All such corrections, which are smooth at $z = 1$, contribute to the moments only by corrections that vanish as powers of N for $N \to \infty$, as indicated. Singular distributions, however, give rise to logarithmic N-dependence, as in

$$\int_0^1 dz\; z^N \left[\frac{\ln^m(1-z)}{1-z}\right]_+ = \frac{(-1)^{m+1}}{m+1}\ln^{m+1} N + \ldots \tag{2.3}$$

Thus, we should think of the large-N behavior as a diagnostic for singular distributions in $1 - z$. Physical cross sections, of course, always involve the inverse transform of the moments.

The functions $\omega_{i\bar{i}\to B}(z, Q/\mu, \alpha_s(\mu^2))$ whose moments appear in Eq. (2.1) are normalized to

$$\omega_{i\bar{i}\to B} = \delta(1-z) + \mathcal{O}(\alpha_s)\,. \tag{2.4}$$

We shall see that the logarithmic N-dependence of moments of ω exponentiates into a useful form.

Before sketching the physical basis and derivation of this result, we observe that, as shown in [31], the same functions $\omega_{i\bar{i}\to B}$ as in Eq. (2.1) contain the singular distributions at $z = 1$ for fixed rapidity,

$$\begin{aligned}\frac{d\sigma_{h_1h_2\to B}}{dQ^2dy} \;=\; & \sum_{i=q,\bar{q},g}\sigma_0\int_\tau^1 dz\int\frac{dx_a}{x_a}\frac{dx_b}{x_b}\,\phi_{i/h_1}(x_a,\mu^2)\;\phi_{\bar{i}/h_2}(x_b,\mu^2) \\ & \times\,\delta\left(z-\frac{Q^2}{x_ax_bS}\right)\;\delta\left(y-\frac{1}{2}\ln\frac{x_a}{x_b}\right)\omega_{i\bar{i}\to B}(z,Q/\mu,\alpha_s(\mu^2))\,.\end{aligned} \tag{2.5}$$

Thus, we may compute singular threshold corrections for the cross section at fixed rapidity from the inclusive cross section.

The resummation of singular distributions in $1-z$ may be carried out in a number of ways, but all are based on the observation that the perturbative function $\omega_{i\bar{i}}$ may be computed in dimensionally regulated perturbation theory by choosing the external hadrons to be *partons*. For us the most informative approach is based on the ability to reexpress moments of the inclusive Drell-Yan cross section Eq. (2.1), with, for instance, $h_1 = q$, $h_2 = \bar{q}$ [7, 8]. In this form, all logarithmic N behavior is factorized into alternate parton distributions ψ, along with a "soft-parton" function U, which describes the exchange and emission of soft gluons by the annihilating quark pair [7],

$$\begin{aligned}\int_0^1 d\tau \tau^{N-1} \frac{1}{\sigma_0}\frac{d\sigma_{i\bar{i}\to B}}{dQ^2}(\tau, Q^2, \epsilon) &= \left|H\left(\frac{Q}{\mu}, \alpha_s(\mu^2)\right)\right|^2 \tilde{U}^{(i)}\left(\frac{Q}{N\mu}, \alpha_s(\mu^2)\right) \\ &\quad \times \tilde{\psi}_{i/i}\left(N, \frac{Q}{\mu}, \epsilon\right)\ \tilde{\psi}_{\bar{i}/\bar{i}}\left(N, \frac{Q}{\mu}, \epsilon\right) \\ &\quad + \mathcal{O}(1/N)\,. \end{aligned} \tag{2.6}$$

This factorization is illustrated in Fig. 1. H is an infrared-safe function that is independent of N. Let us discuss the remaining functions, which absorb the N-dependence, in turn.

The modified parton distributions ψ in Eq. (2.6) are conveniently defined in the partonic center of mass system at fixed *energy*, rather than a light-like momentum fraction. For $i = q$, for example, we have[2]

$$\psi_{q/q}(x, 2p_0/\mu, \epsilon) = \frac{1}{2\pi 2^{3/2}} \int_{-\infty}^{\infty} dy_0\ \mathrm{e}^{-ixp_0y_0} \langle q(p)|\bar{q}(y_0, \vec{0})\ \frac{1}{2} n\cdot\gamma\ q(0)|q(p)\rangle\,, \tag{2.7}$$

where the matrix element is evaluated in $A_0 = 0$ gauge in the partonic c.m. frame. The vector n is lightlike in the opposite direction from p^μ, so that for $\vec{p}$ in the ± 3 direction, $n\cdot\gamma = \gamma^\mp$. $\psi_{\bar{q}/\bar{q}}$ and $\psi_{g/g}$ may be constructed similarly. Defined in this fashion, ψ absorbs all the collinear divergences of the Drell-Yan cross section in the $\tau \to 1$ limit [7]. Its moments may thus be factorized into a product of moments of the corresponding light-cone parton distribution ϕ, defined in any scheme, times an infrared safe function.

[2]This form includes an explicit spin average. Single-particle states are normalized by $\langle 0|q(0)|p, s\rangle = u(p, s)$. The factor of $1/2^{3/2}$ then normalizes the lowest order to $\delta(1-x)$.



(a)



(b)

Fig. 1. (a) Factorization for the Drell-Yan process near partonic threshold, Eq. (2.6). (b) The soft-gluon function U. The double lines represent ordered exponentials in Eq. (2.10).

The function $U^{(i)}$ represents the coupling of soft gluons to the annihilating partons. So long as these soft gluons are not collinear to the active partons, this coupling is well approximated by ordered exponentials, or Wilson lines, which we denote as

$$\Phi_\beta(\lambda_2,\lambda_1;x) = P\ \exp\left(-ig\int_{\lambda_1}^{\lambda_2} d\lambda'\beta\cdot A(\lambda'\beta+x)\right)\,, \tag{2.8}$$

with A the gauge field in the appropriate representation of the gauge group, and β a velocity four-vector, in the direction of the particle whose interactions with soft gluons are being approximated. In the case of a quark, the gauge fields are in the fundamental representation, for a gluon in the adjoint. The operator P orders group products in the same sense as ordering in the integration variable λ, with the A's with lower values of λ to the right. In this notation, $\Phi_\beta(\lambda_2,\lambda_1;x)$ represents (for example) a quark that propagates from $x+\lambda_1\beta$ to $x+\lambda_2\beta$, while $\Phi_\beta(\lambda_1,\lambda_2;x)$ represents an antiquark with the same propagation. The construction of $U^{(i)}$ from the Φ's is actually a two-step process, which we now describe.

Starting with the Wilson lines, we first construct a (dimensionless) "eikonal singlet cross section"

$$\begin{aligned}\sigma_{\rm DY}^{\rm (eik)}\left((1-w)Q/\mu,\alpha_s(\mu^2),\epsilon\right) &= \frac{Q}{2\pi}\int dx_0\ {\rm e}^{i(1-w)Qx_0}\ \langle 0|\bar{T}\left[W^\dagger_{b_1b_2}(x_0,\vec{0})\right] \\ &\quad\times T\left[W_{b_2b_1}(0)\right]|0\rangle\,,\end{aligned} \tag{2.9}$$

where $W(x)$ is the product of Wilson lines representing the pair, coupled by a color-singlet vertex at the point x,

$$W_{b_2,b_1}(x) = \delta_{a_1a_2}\ \Phi_{\beta_2}(-\infty,0;x)_{b_2a_2}\Phi_{\beta_1}(0,-\infty;x)_{a_1b_1}\,. \tag{2.10}$$

In Eq. (2.9), T represents time-ordering, and $\bar{T}$ anti-time ordering. When dealing with QCD hard-scattering below, we shall generalize this product, by incorporating nontrivial color structure at the vertex between the lines. The quantity $(1-w)Q$ is the energy emitted by the Wilson lines [27, 32], which is the relevant measure of phase space near partonic threshold in the center of mass frame of the colliding partons [7]. In Eq. (2.9), an appropriate normalization is understood, so that at lowest order $\sigma_{\rm DY}^{\rm (eik)} = \delta(1-w)$.

The eikonal cross section represents the emission of soft gluons from the annihilating pair. The scale invariance of the Wilson lines implies, however,

that it contains ultraviolet, in addition to infrared and collinear, divergences, corresponding to the renormalization of the composite operator introduced by the vertex in Eq. (2.10) where the eikonal lines join [23, 24, 25, 26]. A natural "scheme" for this renormalization is specified by demanding that the integral of $\sigma_{\rm DY}^{\rm (eik)}(1-w)$ from $w=0$ to $w=1$ be normalized to unity,

$$\int_0^1 dw\; \sigma_{\rm DY}^{\rm (eik)}\left((1-w)Q/\mu, \alpha_s(\mu^2), \epsilon\right) = 1\,. \tag{2.11}$$

That is, we demand that all corrections to $\sigma_{\rm DY}^{\rm (eik)}$ vanish for the fully inclusive cross section. With this normalization, beyond lowest order $\sigma_{\rm DY}^{\rm (eik)}$ is a sum of "plus" distributions in $1-w$,

$$\begin{aligned}\sigma_{\rm DY}^{\rm (eik)}\left((1-w)Q/\mu, \alpha_s(\mu^2), \epsilon\right) &= \delta(1-w) \\ &\quad + \sum_{n\geq 0} s_n\left(Q/\mu, \alpha_s(\mu^2), \epsilon\right)\left[\frac{\ln^n(1-w)}{1-w}\right]_+ ,\end{aligned} \tag{2.12}$$

where the coefficients s_n are power series in the coupling.

Even after renormalization, $\sigma_{\rm DY}^{\rm (eik)}$ includes the couplings of fast-moving gluons that are collinear to the incoming Wilson lines. Such gluons, whether virtual or real, are already included in the distributions ψ, and their contributions should be removed from the soft gluon function, $U^{(i)}$, to avoid double counting. This is possible because noncollinear soft gluons factor from collinear gluons through the use of Ward identities [33]. Physically, soft radiation cannot distinguish between physical and eikonal light-like sources. Thus, moments of $\sigma_{\rm DY}^{\rm eik}$ are factorizable in the same manner as the quark-antiquark cross section cross section, Eq. (2.6), into moments of the (identical) soft-parton function U times moments of "jet" functions $\tilde{j}_{1,2}$, analogous to the ψ's, Eq. (2.7), but with the incoming partons represented by products of eikonal lines (in $A_0=0$ gauge) as

$$\begin{aligned} j_1\left(\frac{(1-w_1)Q}{\mu}, \alpha_s(\mu^2), \epsilon\right) &= \frac{Q}{2\pi}\int_{-\infty}^{\infty} dy_0\; {\rm e}^{-i(1-w_1)Qy_0} \\ &\quad \times\langle 0|\;{\rm Tr}\Big\{\;\bar{T}[\Phi^\dagger_{\beta_1}(0,-\infty;y)] \\ &\quad \times T[\Phi_{\beta_1}(0,-\infty;0)]\;\Big\}\;|0\rangle\,, \end{aligned} \tag{2.13}$$

with $y^\nu = (y_0, \vec{0})$ a vector at the spatial origin. The "antiquark" distribution j_2 is defined analogously, but with the Wilson line in the β_2 direction and ordered from 0 to $-\infty$.

In these terms, the eikonal version of the moment relation Eq. (2.6) is

$$\begin{aligned}\tilde{\sigma}_{\rm DY}^{\rm (eik)}\left(Q/N\mu, \alpha_s(\mu^2), \epsilon\right) &= \tilde{U}^{(i)}\left(\frac{Q}{N\mu}, \alpha_s(\mu^2)\right) \\ &\quad \times \tilde{j}_1\left(\frac{Q}{N\mu}, \alpha_s(\mu^2), \epsilon\right)\ \tilde{j}_2\left(\frac{Q}{N\mu}, \alpha_s(\mu^2), \epsilon\right),\end{aligned} \tag{2.14}$$

which may be used to construct the soft function U. This prescription eliminates diagrams that have divergences from gluons collinear to the external eikonal lines of U, a fact which we shall use below. Because the j_i's absorb all collinear divergences, U is an infrared safe sum of plus distributions.

Logarithms of N exponentiate in each of the factors of Eq. (2.6) [7]. As emphasized in Ref. [10], this is a result of the constraint that the product is independent of both the gauge choice and the factorization scale, although its individual factors are not. Such combined constraints organize logarithms associated with both soft and collinear momentum configurations, and are at the basis of Sudakov resummation [34]. Comparing Eq. (2.1) with $h_1 = i$, $h_2 = \bar{i}$ with Eq. (2.6), we can solve for $\tilde{\omega}(N)$, in terms of the calculable N-dependence of ψ, ϕ and $U^{(i)}$ [7],

$$\tilde{\omega}_{i\bar{i}\to B}(N) = \left[\frac{\tilde{\psi}_{i/i}(N,1,\epsilon)}{\tilde{\phi}_{i/i}(N,Q^2,\epsilon)}\right]^2 \left|H\left(\frac{Q}{\mu}, \alpha_s(\mu^2)\right)\right|^2 \tilde{U}^{(i)}\Big(\frac{Q}{N\mu}, \alpha_s(\mu^2)\Big), \tag{2.15}$$

where we have used $\psi_{i/i} = \psi_{\bar{i}/\bar{i}}$, and similarly for the ϕ's, and have suppressed various arguments.

The result of this procedure applied to the Drell-Yan cross section is

$$\begin{aligned}\tilde{\omega}_{i\bar{i}\to B}(N) &= A(\alpha_s(Q^2))\ \exp\left[-\int_0^1 dz \frac{z^{N-1}-1}{1-z}\right. \\ &\quad \left.\times\left(\int_{(1-z)^2}^{(1-z)^{m_S}} \frac{d\lambda}{\lambda} g_1^{(i\bar{i})}[\alpha_s(\lambda Q^2)] + g_2^{(i\bar{i})}[\alpha_s((1-z)^{m_s}Q^2)]\right)\right],\end{aligned} \tag{2.16}$$

where for DIS ($\overline{\rm MS}$) factorization scheme $m_S = 1\,(0)$. $A(\alpha_s)$ is a constant function of the coupling, and the $g_{1,2}$ are finite functions of their arguments. $g_1^{(ab)}$ and $g_2^{(ab)}$ are universal among hard cross sections and color structures for given incoming partons a and b, but depend on whether these partons are quarks or gluons. To reach the accuracy of next-to-leading logarithms in the exponents, we need g_1 only to two loops, with leading logarithms coming entirely from its one-loop approximation, and g_2 only to a single loop. More explicitly, we take [13]

$$g_1^{(ab)} = (C_a + C_b)\left(\frac{\alpha_s}{\pi} + \frac{1}{2}K\left(\frac{\alpha_s}{\pi}\right)^2\right), \tag{2.17}$$

with $C_i = C_F$ (C_A) for an incoming quark (gluon), and with K given by

$$K = C_A\left(\frac{67}{18} - \frac{\pi^2}{6}\right) - \frac{5}{9}n_f\,, \tag{2.18}$$

where n_f is the number of quark flavors. g_2 is given for quarks in the DIS scheme by

$$g_2^{(q\bar{q})} = -\frac{3}{2}C_F\,\frac{\alpha_s}{\pi}\,, \tag{2.19}$$

and it vanishes in the $\overline{\rm MS}$ scheme. The resummed N-dependence thus depends on the factorization scheme, and is quite different between DIS and $\overline{\rm MS}$ schemes [8, 10]. This difference is in principle compensated for by differences in the N-dependence of the distributions themselves.

The leading logarithms in the exponent of Eq. (2.16) are from the ratio of ψ to ϕ in Eq. (2.15). The function $U^{(i)}$ contributes at next-to-leading logarithm. To achieve this accuracy we need only a one-loop calculation, and we find [7]

$$\begin{aligned}\tilde{U}^{(i)}(N) &= \exp\left[\int_0^1 dz\frac{z^{N-1}-1}{1-z}\nu^{(i)}(\alpha_s((1-z)^2Q^2))\right]\\ &= \exp\left[\int_0^1 dz\frac{z^{N-1}-1}{1-z}\left(\int_{(1-z)^2}^{(1-z)^{m_S}}\frac{d\lambda}{\lambda}\beta(g(\lambda Q^2))\nu^{(i)}[\alpha_s(\lambda Q^2)]\right.\right.\\ &\qquad \left.\left.+\nu^{(i)}[\alpha_s((1-z)^{m_s}Q^2)]\right)\right]\,,\end{aligned} \tag{2.20}$$

with $\nu^{(q,g)} = 2C_{F,A}(\alpha_s/\pi) + \ldots$ In the second form, $\beta(g)$ is the QCD beta function, and we have followed [9] in reexpressing $U^{(i)}$ into the form of Eq.

(2.16). In this connection, we note that in axial gauges $U^{(i)}$ receives noncancelling contributions only from the "gauge" term $k^\mu\, k^\nu/(n\cdot k)^2$ in the gluon propagator. At least to this order, it is natural to use Ward identities to factor such terms into the parton distributions ψ.

2.2 Resummation for QCD Hard Scattering

Generalizing Eq. (2.5), we consider the production of a pair of heavy quarks (or jets) at rapidity y, and at scattering angle θ in the pair center of mass frame, as

$$\begin{aligned}\frac{d\sigma_{h_1h_2\to Q\bar{Q}}}{dQ^2dyd\cos\theta} &= \sum_{f=q,\bar{q},g}\int_\tau^1 dz\int\frac{dx_a}{x_a}\frac{dx_b}{x_b}\,\phi_{f/h_1}(x_a,\mu^2)\;\phi_{\bar{f}/h_2}(x_b,\mu^2)\\ &\quad\times\delta\left(z-\frac{Q^2}{x_ax_bS}\right)\,\delta\left(y-\frac{1}{2}\ln\frac{x_a}{x_b}\right)\\ &\quad\times\hat{\sigma}_{f\bar{f}\to Q\bar{Q}}\left(z,\frac{Q}{\mu},Q,\theta,\alpha_s(\mu^2)\right)\\ &\quad+\ldots\,.\end{aligned}\tag{2.21}$$

Corrections, due to other partonic flavor combinations, are less singular by a power of $1-z$ as $z\to 1$ in the hard scattering, and their moments are suppressed by powers of $1/N$, as in Eq. (2.1). For definiteness, we shall concentrate on heavy quark production, but our considerations are more general. As in the case of Drell-Yan cross sections, the singular behavior of the hard-scattering functions $\hat{\sigma}$ (the analog of $\omega_{q\bar{q}}$ above) at fixed rapidity may be computed from the inclusive partonic cross section [31]. Following Ref. [10], we begin with the factorization properties of heavy quark production and other QCD hard-scattering processes in the elastic limit. We shall see that these depend upon the directions of the outgoing heavy quarks in the partonic center of mass frame. As a result, resummation must be carried out at fixed directions for the heavy quarks. The reasoning is then very similar to that for the Drell-Yan process discussed above. Note in particular, that the partonic subreaction remains an annihilation (or gluon fusion) process.

To apply our methods to heavy quark and other QCD production mechanisms, Eq. (2.6) must be generalized in a number of ways. First, the underlying hard process involves not only a singlet color configuration, but any color tensor that may be constructed from the color representations of the

incoming partons. Second, the outgoing heavy quarks (or other particles), as well as the incoming light partons, act as sources of soft gluons. Our analysis is simplified, however, by the following observation. Near partonic threshold, the emission of light quarks into the final state is suppressed by factors of $1-z$, aside from pairs produced by soft gluons. As a result, although different color structures may "mix" due to soft gluon emission, the *flavor* structure of the hard scattering is unchanged by all corrections that are singular in $1-z$. Thus in the case of heavy quark production, we may analyze light quark annihilation and gluon fusion separately.

To take these new features into account, we generalize the Drell-Yan factorization to QCD hard scattering initiated by light partons f and $\bar{f}$ with the following expressions, which are the analogs of Eqs. (2.1) (for $h_1 = f$, $h_2 = \bar{f}$) and (2.6),

$$\begin{aligned}\int_0^1 d\tau \tau^{N-1} &\frac{d\sigma_{f\bar{f}\to Q\bar{Q}}}{dQ^2}(\tau, Q^2, \epsilon) \\ &= \tilde{\phi}_{f/f}(N, \mu^2, \epsilon)\, \hat{\sigma}_{f\bar{f}\to Q\bar{Q}}(N, Q/\mu, Q, \alpha_s(\mu^2))\, \tilde{\phi}_{\bar{f}/\bar{f}}(N, \mu^2, \epsilon) \\ &= \sum_{IJ} h_J^*\left(\frac{Q}{\mu}, \alpha_s(\mu^2)\right)\, \tilde{S}_{JI}\Big(\frac{Q}{\mu N}, \alpha_s(\mu^2)\Big)\, h_I\left(\frac{Q}{\mu}, \alpha_s(\mu^2)\right) \\ &\quad \times \tilde{\psi}_{f/f}\left(N, \frac{Q}{\mu}, \epsilon\right)\, \tilde{\psi}_{\bar{f}/\bar{f}}\left(N, \frac{Q}{\mu}, \epsilon\right) + \mathcal{O}(1/N)\,. \end{aligned} \tag{2.22}$$

This factorization is illustrated in Fig. 2, for the process

$$f(p_a) + \bar{f}(p_b) \to \bar{Q}(p_1) + Q(p_2)\,, \tag{2.23}$$

in which $f\bar{f}$ represents a pair of light quarks or gluons that annihilate or fuse, respectively, into a heavy quark pair.

This factorization differs from Eq. (2.6) only in taking into account color exchange at the hard scatterings, h_I in the amplitude, and h_J^* in the complex conjugate. For simplicity, we omit the partonic indices $f, \bar{f}$ in h_I and h_J^*, although these functions certainly depend on incoming flavor. For quark-antiquark annihilation, the indices I and J range over two values, corresponding to annihilation in a singlet or octet in the s-channel (or t channel). For other hard-scatterings, such as gluon-gluon, the same pattern applies, but with I and J ranging over color tensors appropriate to possible combinations of gluonic representations in the initial state with (heavy) quark representations in the final state.



(a)



(b)

Fig. 2. (a) Factorization for heavy quark production near partonic threshold, Eq. (2.22). (b) The soft-gluon function S_{JI}, in which the vertices T link ordered exponentials as in Eq. (2.24).

The color tensor of the hard scattering serves to tie together the ordered exponentials to which soft gluons couple, generalizing the singlet vertex joining the eikonal lines in Eq. (2.10) above,

$$\begin{aligned} w_I(x)_{\{c_k\}} &= \Phi_{v_j}(\infty,0;x)_{c_j,d_j}\,\Phi_{v_i}(0,\infty;x)_{d_i,c_i} \\ &\quad \times (T_I)_{d_b d_a, d_j d_i}\;\Phi_{v_a}(-\infty,0;x)_{c_a,d_a}\Phi_{v_b}(0,-\infty;x)_{d_b,c_b}\,, \end{aligned} \tag{2.24}$$

where Φ_{v_a} is in the color representation flavor f, with velocity v_a. Notice that only the Wilson lines of the incoming partons have light-like velocity vectors. Next, by analogy to Eq. (2.9) for the eikonal cross section in Drell-Yan, we define,

$$\begin{aligned} \sigma_{JI}^{(\mathrm{eik})}\left((1-w)Q/\mu,\alpha_s(\mu^2)\right) &= \frac{Q}{2\pi}\int dx_0\;\mathrm{e}^{i(1-w)Qx_0}\;\langle 0|\bar{T}\left[w_J^\dagger(x_0,\vec{0})_{\{b_i\}}\right] \\ &\quad \times T\left[w_I(0)_{\{b_i\}}\right]|0\rangle\,. \end{aligned} \tag{2.25}$$

As in the Drell-Yan case, this cross section includes unphysical collinear divergences associated with its incoming, lightlike eikonal lines. These are eliminated in the soft function S_{JI} by refactorizing $\sigma_{JI}^{(\mathrm{eik})}$ in the same manner as $\sigma_{\mathrm{DY}}^{(\mathrm{eik})}$, Eq. (2.14), through moments,

$$\tilde{\sigma}_{JI}^{(\mathrm{eik})}(N) = \tilde{S}_{JI}(N)\,\tilde{j}_1(N)\,\tilde{j}_2(N)\,. \tag{2.26}$$

Now, comparing Eqs. (2.6) and (2.22), we see that the N-dependence of the cross section for heavy-quark production from a set of initial-state partons is given by the following generalization of Eq. (2.15),

$$\hat{\sigma}_{f\bar{f}\to Q\bar{Q}}(N) = \left[\frac{\tilde{\psi}_{f/f}(N,1,\epsilon)}{\tilde{\phi}_{f/f}(N,Q^2,\epsilon)}\right]^2 \sum_{IJ} h_J^*(Q/\mu)\;\tilde{S}_{JI}(Q/N\mu)\;h_I(Q/\mu)\,, \tag{2.27}$$

where the sum over I and J is, as usual, over color tensors. To determine the singular behavior at partonic threshold, we only need to resum logarithmic N-dependence in each S_{JI}.

The soft matrix S_{JI} depends on N through the ratio $Q/(N\mu)$. Its N-dependence, then, can be resummed by renormalization group analysis [22, 24, 25, 26, 35]. The composite operator in Eq. (2.24), which defines S_{JI}, requires renormalization. Nevertheless, the product $h_J^* S_{JI} h_I$ of the soft function and the hard factors needs no overall renormalization, because the UV

divergences of S_{JI} associated with the new composite operator are balanced by those of the h's. This situation is summarized by the conditions [24, 10]

$$\begin{aligned} h_I^{(0)} &= \prod_{i=1}^{2} Z_i^{-1/2} \left(Z_S^{-1}\right)_{IC} h_C \\ h^{*(0)}_J &= \prod_{i=1}^{2} Z_i^{-1/2} \, h^*_D \, [(Z_S^\dagger)^{-1}]_{DJ} \\ S_{JI}^{(0)} &= (Z_S^\dagger)_{JB} S_{BA} Z_{S,AI}, \end{aligned} \tag{2.28}$$

where $h^{(0)}$ and $S^{(0)}$ denote the unrenormalized quantities, Z_i is the renormalization constant of the ith incoming partonic field external to h_I and $Z_{S,IJ}$ is a matrix of renormalization constants, which describe the renormalization of the soft function, including mixing of color structures. Z_S is defined to include the wave function renormalization necessary for the outgoing eikonal lines that represent the heavy quarks.

From Eq. (2.28), the soft function S_{JI} satisfies the renormalization group equation [24]

$$\left(\mu\frac{\partial}{\partial\mu} + \beta(g)\frac{\partial}{\partial g}\right) S_{JI} = -(\Gamma_S^\dagger)_{JB} S_{BI} - S_{JA}(\Gamma_S)_{AI}\,, \tag{2.29}$$

where the anomalous dimension matrices may be found by explicit renormalization of the soft function. Following [24], we find it convenient to compute the matrix in a minimal subtraction scheme. To be explicit, taking $\epsilon = 4-n$ the matrix of anomalous dimensions may be found at one loop from the matrix of renormalization constants by

$$\Gamma_S(g) = -\frac{g}{2}\frac{\partial}{\partial g}\mathrm{Res}_{\epsilon\to 0} Z_S(g,\epsilon). \tag{2.30}$$

The determination of the anomalous dimension matrices Γ_S will be the subject of the following sections, for heavy quark production through the annihilation of light quarks and gluon fusion.

The solution to the renormalization group equation (2.29) organizes logarithms of N in the soft function. In general, this matrix equation involves a scale-dependent mixing of tensors, and has no closed expression. At the level of leading logarithms of N in S_{JI}, and therefore at next-to-leading logarithm of N in the cross section as a whole, we may simplify by choosing a color

basis in which the anomalous dimension matrix is diagonal, with eigenvalues λ_I for each basis color tensor labelled by I. At this level of approximation, we have

$$\tilde{S}_{JI}\left(\frac{Q}{N\mu_2},\,\alpha_s(\mu_2^2)\right) \;=\; \tilde{S}_{JI}\left(\frac{Q}{N\mu_1},\,\alpha_s\left(\mu_1^2\right)\right) \times\exp\left[-\int_{\mu_1}^{\mu_2}\frac{d\bar{\mu}}{\bar{\mu}}[\lambda_I(\alpha_s(\bar{\mu}^2))+\lambda_J^*(\alpha_s(\bar{\mu}^2))]\right]. \tag{2.31}$$

The coherent effects of soft gluon emission are all contained in this expression, which we now employ to determine $\hat{\sigma}_{f\bar{f}\to Q\bar{Q}}$, Eq. (2.27), to NLL accuracy.

Our first step is to rewrite Eq. (2.27) slightly, to isolate differences and similarities to the Drell-Yan result of Eq. (2.15),

$$\hat{\sigma}_{f\bar{f}\to Q\bar{Q}}(N) \;=\; \left[\frac{\tilde{\psi}_{f/f}(N,1,\epsilon)}{\tilde{\phi}_{f/f}(N,Q^2,\epsilon)}\right]^2 \tilde{U}^{(f)}(Q/N\mu) \times\frac{\sum_{IJ}h_J^*(Q/\mu)\;\tilde{S}_{JI}(Q/N\mu)\;h_I(Q/\mu)}{\tilde{U}^{(f)}(Q/N\mu)}\,. \tag{2.32}$$

We have simply multiplied and divided by $U^{(f)}(Q/N\mu)$, defined in Eq. (2.14). The first factor in Eq. (2.32) is just the resummed singlet cross section (2.16). The new factor is to be treated using Eq. (2.31) and the explicit expression of U, Eq. (2.20).

Combining the results just discussed, and picking a color basis in which moments with respect to z exponentiate, we find, at NLL accuracy

$$\begin{aligned}&\hat{\sigma}_{f\bar{f}\to Q\bar{Q}}(N)\\ &\quad= A'(\alpha_s(Q^2))\;h_J^*\left(1,\alpha_s(Q^2)\right)\;\tilde{S}_{JI}\Big(1,\alpha_s([Q/N]^2)\Big)\;h_I\left(1,\alpha_s(Q^2)\right)\\ &\qquad\times\exp\Big[E_{JI}(N,\theta,Q^2)\Big]\,,\end{aligned} \tag{2.33}$$

where A' is an overall constant and where the color-dependent exponents are given by

$$E_{JI}(N,\theta,Q^2) \;=\; -\int_0^1 dz\frac{z^{N-1}-1}{1-z}\Big[\int_{(1-z)^2}^{(1-z)^{m_S}}\frac{d\lambda}{\lambda}g_1^{(ab)}[\alpha_s(\lambda Q^2)]$$

$$+ g_2^{(ab)}[\alpha_s((1-z)^{m_S}Q^2)]$$
$$+ g_3^{(I)}[\alpha_s((1-z)^2Q^2),\theta] + g_3^{(J)*}[\alpha_s((1-z)^2Q^2),\theta]\Big] . \tag{2.34}$$

$g_3^{(I)}$ summarizes soft logarithms that depend directly on color exchange in the hard scattering, and hence also on the identities and relative directions of the colliding partons (through θ), both incoming and outgoing. As pointed out above, one-loop contributions to g_3 may always be absorbed into the one-loop contribution to g_2 and the two-loop contribution to g_1 [9]. Because $g_3^{(I)}$ depends upon I, however, it is advantageous to keep this nonfactoring color-dependence separate.

Given a choice of incoming and outgoing partons, next-to-leading logarithms in the moment variable N exponentiate as in (2.33) in the color tensor basis that diagonalizes $\Gamma_{S,JI}^{(ab\to cd)}$, with eigenvalues λ_I. The resulting soft gluon anomalous dimension $g_3^{(I)}$ is then simply

$$g_3^{(I)}[\alpha_s,\theta] = -\lambda_I[\alpha_s,\theta] + \frac{1}{2}\nu^{(i)}(\alpha_s)\,, \tag{2.35}$$

where the eigenvalues are complex in general, and depend on the relative directions of the incoming and outgoing partons, as shown.

3 Applications to $q\bar{q} \to Q\bar{Q}$

These considerations may be illustrated by heavy quark production through light quark annihilation,

$$q(p_a) + \bar{q}(p_b) \to \bar{Q}(p_1) + Q(p_2)\,. \tag{3.1}$$

In this case, as in quark-quark elastic scattering [24, 26], the anomalous dimension matrix is two-dimensional. Following Ref. [36] we define the invariants

$$s = (p_a + p_b)^2\,, \quad t_1 = (p_a - p_1)^2 - m^2\,, \quad u_1 = (p_b - p_1)^2 - m^2\,, \tag{3.2}$$

with m the heavy quark mass, which satisfy

$$s + t_1 + u_1 = 0 \tag{3.3}$$

at partonic threshold. We also define dimensionless vectors v_i^μ by

$$p_i^\mu = \mathcal{Q} v_i^\mu \tag{3.4}$$

which obey $v_i^2 = 0$ for the light incoming quarks and $v_i^2 = m^2/\mathcal{Q}^2$ for the outgoing heavy quarks. Note that $\mathcal{Q}$ satisfies the kinematic relation $s = 2\mathcal{Q}^2$.

We now calculate the matrix $\Gamma_S(g)$ for this case. Renormalization associated with incoming light partons is factorized into the wave functions ψ of the previous section. We therefore need consider only diagrams which are not part of the eikonal "jets" j_i, as in Eq. (2.13). In particular, we may omit the UV structure of the incoming light-like eikonal lines.

The UV divergent $O(\alpha_s)$ contribution to each S_{JI} is the sum of graphs illustrated by Fig. 3. Most interesting are the "vertex corrections" in Fig. 3(a). The counterterms for S are the ultraviolet divergent coefficients times our basis color tensors, denoted c_i

$$\begin{aligned} S_1 &= c_1 Z_{S,11} + c_2 Z_{S,21}, \\ S_2 &= c_1 Z_{S,12} + c_2 Z_{S,22}\,, \end{aligned} \tag{3.5}$$

where we will specify a convenient basis for the c_i below.

In our calculations we employ the general axial gauge gluon propagator,

$$D^{\mu\nu}(k) = \frac{-i}{k^2 + i\epsilon} N^{\mu\nu}(k), \quad N^{\mu\nu}(k) = g^{\mu\nu} - \frac{n^\mu k^\nu + k^\mu n^\nu}{n\cdot k} + n^2 \frac{k^\mu k^\nu}{(n\cdot k)^2}, \tag{3.6}$$

with n^μ the gauge vector, and eikonal rules for all external lines (Fig. 4). We now review the notation of Ref. [24], which enables us to summarize results for the diagrams of Fig. 3 in a unified manner.

If we denote a typical one-loop correction to c_I as $\omega^{(I)}(\delta_i v_i, \delta_j v_j, \Delta_i, \Delta_j)$, where $\delta = +1\,(-1)$ for the gluon momenta flowing in the same (opposite) direction to the momentum of v_i and $\Delta = +1\,(-1)$ for v_i corresponding to a quark (antiquark), then we have:

$$\begin{aligned} \omega^{(I)} &= c_I\, g^2 \int \frac{d^n q}{(2\pi)^n} \frac{-i}{q^2 + i\epsilon} \left\{ \frac{\Delta_i \Delta_j v_i \cdot v_j}{(\delta_i v_i \cdot q + i\epsilon)(\delta_j v_j \cdot q + i\epsilon)} \right. \\ &\left. - \frac{\Delta_i v_i \cdot n}{(\delta_i v_i \cdot q + i\epsilon)} \frac{P}{(n\cdot q)} - \frac{\Delta_j v_j \cdot n}{(\delta_j v_j \cdot q + i\epsilon)} \frac{P}{(n\cdot q)} + n^2 \frac{P}{(n\cdot q)^2} \right\}, \end{aligned} \tag{3.7}$$



(a)



(b)

Fig. 3. One-loop corrections to S_{JI} for heavy quark production through light quark annihilation: (a) vertex corrections; (b) heavy quark self-energy graphs.



Fig. 4. Feynman rules for eikonal lines in $q\bar{q} \to Q\bar{Q}$. The gluon momentum flows out of the eikonal lines. Group matrices at the vertices are the same as for quark lines.

where c_I is a color tensor. P stands for principal value,

$$\frac{P}{(q\cdot n)^\beta} = \frac{1}{2}\left(\frac{1}{(q\cdot n + i\epsilon)^\beta} + (-1)^\beta \frac{1}{(-q\cdot n + i\epsilon)^\beta}\right). \tag{3.8}$$

We rewrite (3.7) as

$$\begin{aligned}\omega^{(I)} &= c_I\, \mathcal{S}_{ij}\left[I_1(\delta_i v_i, \delta_j v_j) - \frac{1}{2}I_2(\delta_i v_i, n) - \frac{1}{2}I_2(\delta_i v_i, -n)\right. \\ &\quad \left. -\frac{1}{2}I_3(\delta_j v_j, n) - \frac{1}{2}I_3(\delta_j v_j, -n) + I_4(n^2)\right], \end{aligned}\tag{3.9}$$

where the overall sign is given by

$$\mathcal{S}_{ij} = \Delta_i\, \Delta_j\, \delta_i\, \delta_j. \tag{3.10}$$

We now evaluate the ultraviolet poles of the integrals. For the integrals when both v_i and v_j refer to massive quarks we have (with $\epsilon = 4 - n$)

$$\begin{aligned} I_1^{UV\ pole} &= \frac{\alpha_s}{\pi}\frac{1}{\epsilon}L_\beta\,, \\ I_2^{UV\ pole} &= -\frac{\alpha_s}{\pi}\frac{1}{\epsilon}L_i\,, \end{aligned}$$

$$
\begin{aligned}
I_3^{UV\ pole} &= -\frac{\alpha_s}{\pi}\frac{1}{\epsilon}L_j\,, \\
I_4^{UV\ pole} &= -\frac{\alpha_s}{\pi}\frac{1}{\epsilon}\,,
\end{aligned}
\tag{3.11}
$$

where the L_β is the familiar velocity-dependent eikonal function

$$
L_\beta = \frac{1-2m^2/s}{\beta}\left(\ln\frac{1-\beta}{1+\beta}+\pi i\right)\,, \tag{3.12}
$$

with $\beta=\sqrt{1-4m^2/s}$. The L_i and L_j are rather complicated functions of the gauge vector n. We will see shortly that their contributions are cancelled by the inclusion of self energies. Their explicit expressions are:

$$
L_i = \frac{1}{2}[L_i(+n)+L_i(-n)]\,, \tag{3.13}
$$

where

$$
\begin{aligned}
L_i(\pm n) &= \frac{1}{2}\frac{|v_i\cdot n|}{\sqrt{(v_i\cdot n)^2-2m^2n^2/s}} \\
&\quad\left[\ln\left(\frac{\delta(\pm n)\,2m^2/s-|v_i\cdot n|-\sqrt{(v_i\cdot n)^2-2m^2n^2/s}}{\delta(\pm n)\,2m^2/s-|v_i\cdot n|+\sqrt{(v_i\cdot n)^2-2m^2n^2/s}}\right)\right. \\
&\quad\left.+\ln\left(\frac{\delta(\pm n)\,n^2-|v_i\cdot n|-\sqrt{(v_i\cdot n)^2-2m^2n^2/s}}{\delta(\pm n)\,n^2-|v_i\cdot n|+\sqrt{(v_i\cdot n)^2-2m^2n^2/s}}\right)\right]
\end{aligned}
\tag{3.14}
$$

with $\delta(n)\equiv|v_i\cdot n|/(v_i\cdot n)$.

When v_i refers to a massive quark and v_j to a massless quark we have

$$
\begin{aligned}
I_1^{UV\ pole} &= \frac{\alpha_s}{2\pi}\left\{\frac{2}{\epsilon^2}-\frac{1}{\epsilon}\left[\gamma+\ln\left(\frac{v_{ij}^2 s}{2m^2}\right)-\ln(4\pi)\right]\right\}\,, \\
I_2^{UV\ pole} &= -\frac{\alpha_s}{\pi}\frac{1}{\epsilon}L_i\,, \\
I_3^{UV\ pole} &= \frac{\alpha_s}{2\pi}\left\{\frac{2}{\epsilon^2}-\frac{1}{\epsilon}\left[\gamma+\ln(\nu_j)-\ln(4\pi)\right]\right\}\,, \\
I_4^{UV\ pole} &= -\frac{\alpha_s}{\pi}\frac{1}{\epsilon}\,,
\end{aligned}
\tag{3.15}
$$

where

$$\nu_a = \frac{(v_a \cdot n)^2}{|n|^2}\,, \tag{3.16}$$

and $v_{ij} = v_i \cdot v_j$. Note that the double poles cancel in the sum over the I's in Eq. (3.9).

Finally, when both v_i and v_j refer to massless quarks we have [24]

$$\begin{aligned}
I_1^{UV\ pole} &= \frac{\alpha_s}{\pi}\left\{\frac{2}{\epsilon^2} - \frac{1}{\epsilon}\left[\gamma + \ln\left(\delta_i\delta_j\,\frac{v_{ij}}{2}\right) - \ln(4\pi)\right]\right\}\,,\\
I_2^{UV\ pole} &= \frac{\alpha_s}{2\pi}\left\{\frac{2}{\epsilon^2} - \frac{1}{\epsilon}\left[\gamma + \ln(\nu_i) - \ln(4\pi)\right]\right\}\,,\\
I_3^{UV\ pole} &= \frac{\alpha_s}{2\pi}\left\{\frac{2}{\epsilon^2} - \frac{1}{\epsilon}\left[\gamma + \ln(\nu_j) - \ln(4\pi)\right]\right\}\,,\\
I_4^{UV\ pole} &= -\frac{\alpha_s}{\pi}\frac{1}{\epsilon}\,.
\end{aligned} \tag{3.17}$$

Again, note that the double poles cancel.

Our calculations are most easily carried out in a color tensor basis consisting of singlet exchange in the s and u channels,

$$c_1 = \delta_{ab}\delta_{12}\,, \qquad c_2 = \delta_{a2}\delta_{b1}\,. \tag{3.18}$$

The color indices for the incoming (light) quark and antiquark are a and b, respectively, and for the outgoing (heavy) quark and antiquark 2 and 1, respectively.

In the basis (3.18) we find from Eq. (2.30) that the contributions of the diagrams in Fig. 3(a) to the anomalous dimensions are

$$\begin{aligned}
\hat\Gamma_{S,11} &= \frac{\alpha_s}{\pi}C_F\left[\ln\left(\frac{v_{ab}}{2}\right) - L_\beta - \frac{1}{2}\ln(\nu_a\nu_b) - L_1 - L_2 + 2 - \pi i\right]\\
&\quad -\frac{1}{N}\hat\Gamma_{S,21},\\
\hat\Gamma_{S,21} &= \frac{\alpha_s}{2\pi}\ln\left(\frac{v_{a2}v_{b1}}{v_{a1}v_{b2}}\right),\\
\hat\Gamma_{S,12} &= \frac{\alpha_s}{2\pi}\left[\ln\left(\frac{v_{ab}}{2}\right) - \ln(v_{a1}v_{b2}) - L_\beta + \ln\left(\frac{2m^2}{s}\right) - \pi i\right],\\
\hat\Gamma_{S,22} &= \frac{\alpha_s}{\pi}C_F\left[\ln(v_{a2}v_{b1}) - \ln\left(\frac{2m^2}{s}\right) - \frac{1}{2}\ln(\nu_a\nu_b) - L_1 - L_2 + 2\right]\\
&\quad -\frac{1}{N}\hat\Gamma_{S,12}.
\end{aligned} \tag{3.19}$$

The matrix depends, as expected, on the directions of the Wilson lines, which may be reexpressed in terms of ratios of kinematic invariants for the partonic scattering. It is also gauge-dependent.

We now combine this matrix with the (color-diagonal) contributions from the heavy-quark self-energy graphs in Fig. 3(b). The contribution of the self-energy graphs is

$$\frac{\alpha_s}{\pi} C_F (L_1 + L_2 - 2)\, \delta_{IJ}\,, \tag{3.20}$$

which cancels the gauge-dependent L_i of Eq. (3.13). In addition, recalling Eq. (2.35), we subtract half of the anomalous dimension $\nu^{(i)} = 2(\alpha_s/\pi) C_{F,A}$ associated with the soft gluon function $U^{(i)}$ of the Drell-Yan cross section (see Eq. (2.32)). This shifts our matrix by an additional $-(\alpha_s/\pi) C_F \delta_{IJ}$. Combining these contributions with $\hat{\Gamma}_S$ of Eq. (3.19), the full soft-gluon anomalous dimension matrix becomes

$$\begin{aligned}
\Gamma_{S,11} &= \frac{\alpha_s}{\pi} C_F \left[\ln\left(\frac{v_{ab}}{2}\right) - L_\beta - \frac{1}{2}\ln(\nu_a \nu_b) - 1 - \pi i \right] \\
&\quad -\frac{1}{N}\Gamma_{S,21}, \\
\Gamma_{S,21} &= \frac{\alpha_s}{2\pi} \ln\left(\frac{v_{a2} v_{b1}}{v_{a1} v_{b2}}\right), \\
\Gamma_{S,12} &= \frac{\alpha_s}{2\pi} \left[\ln\left(\frac{v_{ab}}{2}\right) - \ln(v_{a1} v_{b2}) - L_\beta + \ln\left(\frac{2m^2}{s}\right) - \pi i \right], \\
\Gamma_{S,22} &= \frac{\alpha_s}{\pi} C_F \left[\ln(v_{a2} v_{b1}) - \ln\left(\frac{2m^2}{s}\right) - \frac{1}{2}\ln(\nu_a \nu_b) - 1 \right] \\
&\quad -\frac{1}{N}\Gamma_{S,12}\,.
\end{aligned} \tag{3.21}$$

In the $A^0 = 0$ gauge we have $\nu_a = \nu_b = 1/2$ in the partonic center of mass frame. Using the relations $v_{ab} = 1$, $v_{12} = 1 - 2m^2/s$, $v_{a1} = v_{b2} = -t_1/s$ and $v_{1b} = v_{2a} = -u_1/s$, the anomalous dimension matrix becomes in terms of our invariants s, t_1, and u_1 :

$$\begin{aligned}
\Gamma_{S,11} &= \frac{\alpha_s}{\pi} \left\{ C_F \left[\ln\left(\frac{u_1^2}{t_1^2}\right) - L_\beta - 1 - \pi i \right] - \frac{C_A}{2} \ln\left(\frac{u_1^2}{t_1^2}\right) \right\}, \\
\Gamma_{S,21} &= \frac{\alpha_s}{2\pi} \ln\left(\frac{u_1^2}{t_1^2}\right),
\end{aligned}$$

$$\begin{aligned}
\Gamma_{S,12} &= \frac{\alpha_s}{2\pi}\left[\ln\left(\frac{m^2 s}{t_1^2}\right) - L_\beta - \pi i\right], \\
\Gamma_{S,22} &= \frac{\alpha_s}{\pi}\left\{C_F\left[\ln\left(\frac{u_1^2}{t_1^2}\right) - L_\beta - 1 - \pi i\right]\right. \\
&\quad \left. + \frac{C_A}{2}\left[-\ln\left(\frac{m^2 s}{t_1^2}\right) + L_\beta + \pi i\right]\right\}. \qquad (3.22)
\end{aligned}$$

For arbitrary β and fixed scattering angle, we must solve for the relevant diagonal basis of color structure, and determine the eigenvalues. At "absolute" threshold, $\beta = 0$, where the heavy quarks are produced relatively at rest, we find

$$\begin{aligned}
\Gamma_{S,11}^{\beta\to 0} &= -\frac{\alpha_s}{\pi} C_F\left(\pi i + \frac{\pi i}{2\beta}\right), \\
\Gamma_{S,21}^{\beta\to 0} &= 0, \\
\Gamma_{S,12}^{\beta\to 0} &= \frac{\alpha_s}{2\pi}\left(1 - \pi i - \frac{\pi i}{2\beta}\right), \\
\Gamma_{S,22}^{\beta\to 0} &= \frac{\alpha_s}{\pi}\left[-C_F\left(\pi i + \frac{\pi i}{2\beta}\right) - \frac{C_A}{2}\left(1 - \pi i - \frac{\pi i}{2\beta}\right)\right]. \qquad (3.23)
\end{aligned}$$

It is interesting to note that $\Gamma_S^{\beta\to 0}$ is diagonalized [37] in a basis of singlet and octet exchange in the s channel,

$$c_{\rm singlet} = c_1, \qquad c_{\rm octet} = -\frac{1}{2N} c_1 + \frac{c_2}{2}, \qquad (3.24)$$

with eigenvalues, $\lambda_{\rm singlet}^{\beta\to 0} = \Gamma_{S,11}^{\beta\to 0}$ and $\lambda_{\rm octet}^{\beta\to 0} = \Gamma_{S,22}^{\beta\to 0}$.

The general result in this s channel singlet-octet basis (including the contribution of the self-energy graphs, subtracting the $\nu^{(i)}$ contribution as noted above and in a general axial gauge) is:

$$\begin{aligned}
\Gamma_{S,11}^{(1,8)} &= -\frac{\alpha_s}{\pi} C_F[L_\beta + 1 + \pi i + \ln(2\sqrt{\nu_a \nu_b})], \\
\Gamma_{S,21}^{(1,8)} &= \frac{2\alpha_s}{\pi}\ln\left(\frac{u_1}{t_1}\right), \\
\Gamma_{S,12}^{(1,8)} &= \frac{\alpha_s}{\pi}\frac{C_F}{C_A}\ln\left(\frac{u_1}{t_1}\right),
\end{aligned}$$

$$\Gamma_{S,22}^{(1,8)} = \frac{\alpha_s}{\pi}\left\{C_F\left[4\ln\left(\frac{u_1}{t_1}\right) - \ln(2\sqrt{\nu_a\nu_b}) - L_\beta - 1 - \pi i\right] + \frac{C_A}{2}\left[-3\ln\left(\frac{u_1}{t_1}\right) - \ln\left(\frac{m^2 s}{t_1 u_1}\right) + L_\beta + \pi i\right]\right\}. \quad (3.25)$$

Γ_S is also diagonalized in this singlet-octet basis for arbitrary β when the parton-parton c.m. scattering angle is $\theta = 90^\circ$ (where $u_1 = t_1$) with eigenvalues that may be read off from Eq. (3.25).

It is of interest, of course, to compare the one-loop expansion of our results to known one-loop calculations, at the level of NLO. We give our result as a function of z, since the inverse transforms are trivial. The result is proportional to the Born cross section, the one-loop contributions from $g_1^{(q\bar{q})}$ and $g_2^{(q\bar{q})}$, and $\Gamma_{S,22}^{(1,8)}$. In the DIS scheme the result is

$$\hat{\sigma}_{f\bar{f}\to Q\bar{Q}}(1-z,u_1,t_1,s)^{(1)} = \sigma_{\rm Born}\frac{\alpha_s}{\pi}\frac{1}{1-z}\left\{C_F\left[2\ln(1-z) + \frac{3}{2} + 8\ln\left(\frac{u_1}{t_1}\right) - 2 - 2{\rm Re}\, L_\beta + 2\ln\left(\frac{s}{\mu^2}\right)\right] + C_A\left[-3\ln\left(\frac{u_1}{t_1}\right) + {\rm Re}\, L_\beta - \ln\left(\frac{m^2 s}{t_1 u_1}\right)\right]\right\}, \quad (3.26)$$

where we keep only terms that give rise to $\ln N$. The logarithm of s/μ^2 describes the evolution of the parton distributions. This result for the cross section at fixed values of the $Q\bar{Q}$ invariant mass cannot be compared directly to the one-loop results of [36], because the latter is by contrast a single-particle cross section, where the singular behavior is given in terms of the variable s_4,

$$s_4 = (p_2 + k)^2 - m^2 \approx 2p_2\cdot k\,, \quad (3.27)$$

where $k = p_a + p_b - p_1 - p_2$ is the momentum carried away by the gluon. At partonic threshold, both s_4 and $(1-z)$ vanish, but angular integrals over the gluon momentum reflect the difference in phase space between fixed s_4 and fixed $1 - z \approx 2(p_1 + p_2)\cdot k/s$.

Nevertheless, the cross sections are kinematically equivalent in the $\beta \to 0$ limit, where we may make a direct comparison. Near $\beta = 0$, our cross section becomes

$$\hat{\sigma}_{f\bar{f}\to Q\bar{Q}}(1-z,u_1,t_1,s)^{(1)}|_{\beta\to 0} = \sigma_{\rm Born}\frac{\alpha_s}{\pi}\frac{1}{1-z}\left\{C_F\left[2\ln(1-z) + \frac{3}{2}\right.\right.$$

$$+2\ln\left(\frac{4m^2}{\mu^2}\right)\Bigg] - C_A\Bigg\}\,. \quad (3.28)$$

Near $s = 4m^2$, we may identify $2m^2(1-z) = s_4$, and this expression becomes identical to the $\beta \to 0$ limit of eq. (30) of [36]. Even for $\beta > 0$, the two cross sections remain remarkably close, differing only at first nonleading logarithm in the abelian (C_F^2) term, due to the interplay of angular integrals with leading singularities for the differing treatments of phase space.

As for the Drell-Yan cross section, our analysis applies not only to absolute threshold for the production of the heavy quarks ($\beta = 0$), but also to partonic threshold for the production of moving heavy quarks. When β nears unity, however, the anomalous dimensions themselves develop (collinear) singularities associated with the fragmentation of the heavy quarks, which in principle may be factored into nonperturbative fragmentation functions.

Finally, it is easy to check that the anomalous dimension matrix for heavy outgoing quarks Eq. (3.19) reduces in the limit $m \to 0$ to the anomalous dimension matrix for light outgoing quarks, $q(p_a) + \bar{q}(p_b) \to \bar{q}(p_1) + q(p_2)$, which is [24]:

$$\begin{aligned}
\Gamma^{q\bar{q}\to q\bar{q}}_{S,11} &= \frac{\alpha_s}{\pi} C_F \left[\ln\left(\frac{v_{ab}v_{12}}{4}\right) - \frac{1}{2}\ln(\nu_a\nu_b\nu_1\nu_2) + 2 - 2\pi i\right] \\
&\quad -\frac{1}{N}\Gamma^{q\bar{q}\to q\bar{q}}_{S,21}, \\
\Gamma^{q\bar{q}\to q\bar{q}}_{S,21} &= \frac{\alpha_s}{2\pi}\ln\left(\frac{v_{a2}v_{b1}}{v_{a1}v_{b2}}\right), \\
\Gamma^{q\bar{q}\to q\bar{q}}_{S,12} &= \frac{\alpha_s}{2\pi}\left[\ln\left(\frac{v_{ab}v_{12}}{v_{a1}v_{b2}}\right) - 2\pi i\right], \\
\Gamma^{q\bar{q}\to q\bar{q}}_{S,22} &= \frac{\alpha_s}{\pi} C_F \left[\ln\left(\frac{v_{a2}v_{b1}}{4}\right) - \frac{1}{2}\ln(\nu_a\nu_b\nu_1\nu_2) + 2\right] \\
&\quad -\frac{1}{N}\Gamma^{q\bar{q}\to q\bar{q}}_{S,12}\,. \qquad (3.29)
\end{aligned}$$

4 Applications to $gg \to Q\bar{Q}$

In this section we give the results for the anomalous dimension matrix when the incoming partons are gluons and the outgoing partons are heavy quarks

$$g(p_a) + g(p_b) \to \bar{Q}(p_1) + Q(p_2)\,. \qquad (4.1)$$

For the sake of completeness we also give results for the case when the outgoing quarks are light

$$g(p_a) + g(p_b) \to \bar{q}(p_1) + q(p_2)\,. \tag{4.2}$$

In Fig. 5 we show the UV divergent one-loop contributions to S for $gg \to Q\bar{Q}$.

Our analysis is similar to the one in the previous section. We use the same integrals for the calculation of the $\omega^{(I)}$. The eikonal rules for incoming gluons are slightly modified and are given in Fig. 6. We choose the following basis for the color factors:

$$c_1 = \delta^{ab}\,\delta_{21}, \quad c_2 = d^{abc}\,T^c_{21}, \quad c_3 = if^{abc}\,T^c_{21}. \tag{4.3}$$

Again, the counterterms for S are the ultraviolet divergent coefficients times our basis color tensors:

$$\begin{aligned} S_1 &= c_1 Z_{S,11} + c_2 Z_{S,21} + c_3 Z_{S,31}, \\ S_2 &= c_1 Z_{S,12} + c_2 Z_{S,22} + c_3 Z_{S,32}, \\ S_3 &= c_1 Z_{S,13} + c_2 Z_{S,23} + c_3 Z_{S,33}. \end{aligned} \tag{4.4}$$

Our results for the contributions from vertex corrections in Fig. 5(a) to the anomalous dimension matrix when the outgoing quarks are heavy are:

$$\begin{aligned} \hat{\Gamma}_{S,11} &= \frac{\alpha_s}{\pi}\Bigg\{C_F(-L_\beta - L_1 - L_2 + 1) \\ &\qquad + C_A\left[\ln\left(\frac{v_{ab}}{2}\right) - \frac{1}{2}\ln(\nu_a\nu_b) + 1 - \pi i\right]\Bigg\}, \\ \hat{\Gamma}_{S,21} &= 0, \\ \hat{\Gamma}_{S,31} &= \frac{\alpha_s}{\pi}\ln\left(\frac{v_{a2}v_{b1}}{v_{a1}v_{b2}}\right), \\ \hat{\Gamma}_{S,12} &= 0, \\ \hat{\Gamma}_{S,22} &= \frac{\alpha_s}{\pi}\Bigg\{C_F(-L_\beta - L_1 - L_2 + 1) + \frac{C_A}{2}\Bigg[\ln\left(\frac{v_{ab}}{2}\right) \\ &\qquad + \frac{1}{2}\ln(v_{a1}v_{b2}v_{a2}v_{b1}) + L_\beta - \ln\left(\frac{2m^2}{s}\right) - \ln(\nu_a\nu_b) + 2 - \pi i\Bigg]\Bigg\}, \\ \hat{\Gamma}_{S,32} &= \frac{N^2-4}{4N}\hat{\Gamma}_{S,31}, \end{aligned}$$



(a)



(b)

Fig. 5. One-loop corrections to S_{JI} for heavy quark production through gluon fusion: (a) vertex corrections; (b) heavy quark self-energy graphs.



Fig. 6. Feynman rules for eikonal lines in $gg \to Q\bar{Q}$. The gluon momentum flows out of the eikonal lines. Group factors for vertices attached to incoming lines are those of three-gluon vertices, and for outgoing lines those of quark-gluon vertices.

$$\begin{aligned}
\hat{\Gamma}_{S,13} &= \frac{1}{2}\hat{\Gamma}_{S,31}, \\
\hat{\Gamma}_{S,23} &= \frac{C_A}{4}\hat{\Gamma}_{S,31}, \\
\hat{\Gamma}_{S,33} &= \hat{\Gamma}_{S,22}.
\end{aligned} \tag{4.5}$$

Again, the gauge dependence associated with the heavy quarks is eliminated once we include the self-energy graphs in Fig. 5(b). The contribution of the self-energy graphs (in the diagonal elements only) is, as before,

$$\frac{\alpha_s}{\pi} C_F (L_1 + L_2 - 2)\, \delta_{IJ}. \tag{4.6}$$

In analogy to the previous section, we subtract $\nu^{(i)}/2$, which gives an additional $-(\alpha_s/\pi)C_A$ shift in the diagonal elements. In terms of the Mandelstam invariants s, t_1, and u_1, the anomalous dimension matrix becomes (in a general axial gauge):

$$\begin{aligned}
\Gamma_{S,11} &= \frac{\alpha_s}{\pi}\left[-C_F(L_\beta + 1) - C_A\left(\frac{1}{2}\ln\left(4\nu_a\nu_b\right) + \pi i\right)\right], \\
\Gamma_{S,21} &= 0,
\end{aligned}$$

$$
\begin{aligned}
\Gamma_{S,31} &= \frac{\alpha_s}{\pi}\ln\left(\frac{u_1^2}{t_1^2}\right), \\
\Gamma_{S,12} &= 0, \\
\Gamma_{S,22} &= \frac{\alpha_s}{\pi}\left\{-C_F(L_\beta+1)-\frac{C_A}{2}\left[\ln\left(4\nu_a\nu_b\right)-\ln\left(\frac{t_1u_1}{m^2s}\right)-L_\beta+\pi i\right]\right\}, \\
\Gamma_{S,32} &= \frac{N^2-4}{4N}\Gamma_{S,31}, \\
\Gamma_{S,13} &= \frac{1}{2}\Gamma_{S,31}, \\
\Gamma_{S,23} &= \frac{C_A}{4}\Gamma_{S,31}, \\
\Gamma_{S,33} &= \Gamma_{S,22}.
\end{aligned}
\tag{4.7}
$$

At threshold for the production of heavy quarks at rest, the anomalous dimension matrix becomes diagonal. In $A^0=0$ gauge (where $\nu_a=\nu_b=1/2$) its eigenvalues are

$$
\begin{aligned}
\lambda_1^{\beta\to 0} &= \Gamma_{S,11}^{\beta\to 0}=\frac{\alpha_s}{\pi}\left[-C_F\frac{\pi i}{2\beta}-C_A\pi i\right], \\
\lambda_2^{\beta\to 0} &= \Gamma_{S,22}^{\beta\to 0}=\frac{\alpha_s}{\pi}\left[-C_F\frac{\pi i}{2\beta}+\frac{C_A}{2}\left(-1-\pi i+\frac{\pi i}{2\beta}\right)\right], \\
\lambda_3^{\beta\to 0} &= \Gamma_{S,33}^{\beta\to 0}=\lambda_2^{\beta\to 0}.
\end{aligned}
\tag{4.8}
$$

We also note that the matrix is diagonalized at $\theta=90^\circ$ for arbitrary β, with eigenvalues

$$
\begin{aligned}
\lambda_1(\theta=90^\circ) &= \Gamma_{S,11}(\theta=90^\circ)=\frac{\alpha_s}{\pi}[-C_F(L_\beta+1)-C_A\pi i], \\
\lambda_2(\theta=90^\circ) &= \Gamma_{S,22}(\theta=90^\circ)=\frac{\alpha_s}{\pi}\left\{-C_F(L_\beta+1)\right. \\
&\qquad \left.+\frac{C_A}{2}\left[\ln\left(\frac{t_1^2}{m^2s}\right)+L_\beta-\pi i)\right]\right\}, \\
\lambda_3(\theta=90^\circ) &= \Gamma_{S,33}(\theta=90^\circ)=\lambda_2(\theta=90^\circ).
\end{aligned}
\tag{4.9}
$$

Again, it is interesting to compare the one-loop expansion of our results to the one-loop calculations in [36]. In this case, our calculation is complicated by the fact that the color decomposition is not trivial as it was for $q\bar{q}$. We

have to decompose the Born cross section into three terms according to our color tensor basis. After some algebra our result becomes

$$\begin{aligned}
\hat{\sigma}_{gg\to Q\mathrm{Q}}(1-z,u_1,t_1,s)^{(1)} &= \alpha_s^3 \frac{1}{1-z} K_{gg} B_{\mathrm{QED}}(s,t_1,u_1) \left\{ \frac{t_1 u_1}{s^2} \right. \\
&\quad \times \left[-8N^2 C_F C_A \left(\ln(1-z) + \frac{1}{2} \ln\left(\frac{s}{\mu^2}\right) \right) \right. \\
&\quad -2N^2 C_F C_A \ln\left(\frac{t_1 u_1}{m^2 s}\right) \\
&\quad \left. +4N^2 C_F \left(C_F - \frac{C_A}{2} \right) \mathrm{Re}\, L_\beta + 4N^2 C_F^2 \right] \\
&\quad +8N^2 C_F^2 \left(\ln(1-z) + \frac{1}{2} \ln\left(\frac{s}{\mu^2}\right) \right) \\
&\quad +C_F C_A (N^2-2) \ln\left(\frac{t_1\, u_1}{m^2 s}\right) \\
&\quad +2C_F \left(C_F - \frac{C_A}{2} \right) \mathrm{Re}\, L_\beta \\
&\quad \left. -2C_F^2 (N^2-1) \right\} , \qquad (4.10)
\end{aligned}$$

where

$$B_{\mathrm{QED}}(s,t_1,u_1) = \frac{t_1}{u_1} + \frac{u_1}{t_1} + \frac{4m^2 s}{t_1\, u_1} \left(1 - \frac{m^2 s}{t_1\, u_1} \right) , \qquad (4.11)$$

and $K_{gg} = (N^2-1)^{-2}$ is a color average factor. The logarithms of s/μ^2 describe the evolution of the parton distributions.

As we discussed in the previous section, our result cannot be compared directly to the one-loop results of [36], but as $\beta \to 0$ our expression becomes identical to the $\beta \to 0$ limit of the sum of eqs. (36-38) in [36]:

$$\begin{aligned}
\hat{\sigma}_{gg\to Q\bar{Q}}(1-z,u_1,t_1,s)^{(1)}|_{\beta\to 0} &= \alpha_s^3 \frac{1}{1-z} K_{gg} \left\{ 4N^2 C_F (4C_F - C_A) \right. \\
&\quad \times \left[\ln(1-z) + \frac{1}{2} \ln\left(\frac{4m^2}{\mu^2}\right) \right] \\
&\quad \left. +(N^2+2) C_F C_A - 4N^2 C_F^2 \right\} . \qquad (4.12)
\end{aligned}$$

Again, even for $\beta > 0$, the two cross sections remain remarkably close.

Finally, the anomalous dimension matrix for the case when the outgoing quarks are light is given by

$$\begin{aligned}
\Gamma^{gg\to q\bar{q}}_{S,11} &= \frac{\alpha_s}{\pi}\left\{C_F\left[\ln\left(\frac{v_{12}}{2}\right)-\frac{1}{2}\ln(\nu_1\nu_2)+1-\pi i\right]\right. \\
&\quad \left.+C_A\left[\ln\left(\frac{v_{ab}}{2}\right)-\frac{1}{2}\ln(\nu_a\nu_b)+1-\pi i\right]\right\}, \\
\Gamma^{gg\to q\bar{q}}_{S,21} &= 0, \\
\Gamma^{gg\to q\bar{q}}_{S,31} &= \frac{\alpha_s}{\pi}\ln\left(\frac{v_{a2}v_{b1}}{v_{a1}v_{b2}}\right), \\
\Gamma^{gg\to q\bar{q}}_{S,12} &= 0, \\
\Gamma^{gg\to q\bar{q}}_{S,22} &= \frac{\alpha_s}{\pi}\left\{C_F\left[\ln\left(\frac{v_{12}}{2}\right)-\frac{1}{2}\ln(\nu_1\nu_2)+1-\pi i\right]\right. \\
&\quad \left.+C_A\left[\frac{1}{4}\ln(v_{a1}v_{b2}v_{a2}v_{b1})+\frac{1}{2}\ln\left(\frac{v_{ab}}{v_{12}}\right)-\frac{1}{2}\ln(\nu_a\nu_b)-\ln 2+1\right]\right\}, \\
\Gamma^{gg\to q\bar{q}}_{S,32} &= \frac{N^2-4}{4N}\Gamma^{gg\to q\bar{q}}_{S,31}, \\
\Gamma^{gg\to q\bar{q}}_{S,13} &= \frac{1}{2}\Gamma^{gg\to q\bar{q}}_{S,31}, \\
\Gamma^{gg\to q\bar{q}}_{S,23} &= \frac{C_A}{4}\Gamma^{gg\to q\bar{q}}_{S,31}, \\
\Gamma^{gg\to q\bar{q}}_{S,33} &= \Gamma^{gg\to q\bar{q}}_{S,22}. \qquad (4.13)
\end{aligned}$$

We note that this matrix is also diagonalized at threshold and at $\theta = 90^\circ$, and that the anomalous dimension matrix for heavy outgoing quarks, Eq. (4.5) reduces to the anomalous dimension matrix for light outgoing quarks, Eq. (4.13) in the limit $m \to 0$.

5 Conclusions

We have illustrated the application of a general method for resumming next-to-leading logarithms at the edge of phase space in QCD hard-scattering cross sections. We have given explicit results for singular distributions at partonic threshold in heavy quark production through light quark annihilation and gluon fusion, and for light quark production through gluon fusion. Possible extensions include, of course, dijet and multijet production,

and dijet or heavy quark transverse momentum distributions. Recent estimates of the phenomenological importance of these nonleading terms to heavy quark production indicate that their contribution can be substantial [19]. In addition, an understanding of the all-orders perturbative cross section in QCD processes may shed light on power corrections in hadronic scattering [13, 27, 32, 38]. We hope that our method will be of use in improving the reliability of perturbative QCD calculations for hard scattering.

Acknowledgements

This work was supported in part by the National Science Foundation, under grant PHY9309888 and by the PPARC under grant GR/K54601. We wish to thank Lyndon Alvero, Harry Contopanagos, Gregory Korchemsky, Eric Laenen, and Jack Smith for many helpful conversations.

References

[1] G. Altarelli, R.K. Ellis, and G. Martinelli, Nucl. Phys. B157, 461 (1979);
B. Humpert and W.L. van Neerven, Phys. Lett. 84B, 327 (1979);
J. Kubar-André and F.E. Paige, Phys. Rev. D 19, 221 (1979);
K. Harada, T. Kaneko, and N. Sakai, Nucl. Phys. B155, 169 (1979); B165, 545(E) (1980).

[2] P. Aurenche, A. Douiri, R. Baier, M. Fontannaz, and D. Schiff, Phys. Lett. 140B, 87 (1984);
H. Baer, J. Ohnemus, and J.F. Owens, Phys. Rev. D 42, 61 (1990);
P. Aurenche, P. Chiappetta, M. Fontannaz, J.P. Guillet, and E. Pilon, Nucl. Phys. B399, 34 (1993);
L.E. Gordon and W. Vogelsang, Phys. Rev. D 48, 3136 (1993); D 50, 1901 (1994);
M. Glúck, L.E. Gordon, E. Reya, and W. Vogelsang, Phys. Rev. Lett. 73, 388 (1994);
E.L. Berger, X.-f. Guo, and J. Qiu, Phys. Rev. Lett. 76, 2234 (1996).

[3] P. Nason, S. Dawson, and R.K. Ellis, Nucl. Phys. B303, 607 (1988);
W. Beenakker, H. Kuijf, W.L. van Neerven, and J. Smith Phys. Rev. D

40, 54 (1989);
W. Beenakker, W.L. van Neerven, R. Meng, G.A. Schuler, and J. Smith, Nucl. Phys. B351, 507 (1991);
M. Buza, Y. Matiounine, J. Smith, and W.L. van Neerven, Nucl. Phys. B485, 420 (1997).

[4] F. Aversa, P. Chiappetta, M. Greco, and J.-Ph. Guillet, Z. Phys. C46, 253 (1990);
F. Aversa, P. Chiappetta, L. Gonzales, M. Greco, and J.-Ph. Guillet, Z. Phys. C49, 459 (1991);
S.D. Ellis, Z. Kunszt, and D.E. Soper, Phys. Rev. D 40, 2188 (1989); Phys. Rev. Lett. 69, 1496 (1992);
Z. Kunszt and D.E. Soper, Phys. Rev. D 46, 192 (1992);
W.T. Giele, E.W.N. Glover, and D.A. Kosower, Phys. Rev. D 52, 1486 (1995);
E. Mirkes and D. Zeppenfeld, Phys. Lett. B 380, 205 (1996);
S. Catani and M.H. Seymour, Nucl. Phys. B485, 291 (1997);
M. Klasen and G. Kramer, DESY-97-002, hep-ph/9701247, 1997.

[5] L.R. Surguladze and M.A. Samuel, Rev. Mod. Phys. 68, 259 (1996).

[6] R. Hamberg and W.L. van Neerven, Nucl. Phys. B379, 143 (1992);
E.B. Zijlstra and W.L. van Neerven, Nucl. Phys. B383, 525 (1992); B417, 61 (1994); B426, 245(E) (1994);
S.A. Larin, T. van Ritbergen, and J.A.M. Vermaseren, Nucl. Phys. B427, 41 (1994);
T. van Ritbergen, J.A.M. Vermaseren, S.A. Larin, and P. Nogueira, Int. J. Mod. Phys. C6, 513 (1995);
R. Mertig and W.L. van Neerven, Z. Phys. C70, 637 (1996).

[7] G. Sterman, Nucl. Phys. B281, 310 (1987).

[8] S. Catani and L. Trentadue, Nucl. Phys. B327, 323 (1989).

[9] S. Catani and L. Trentadue, Nucl. Phys. B353, 183 (1991).

[10] H. Contopanagos, E. Laenen, and G. Sterman, Nucl. Phys. B484, 303 (1997).

[11] G. Parisi, Phys. Lett. 90B, 295 (1980);
G. Curci and M. Greco, Phys. Lett. 92B, 175 (1980);
P. Chiappetta, T. Grandou, M. Le Bellac, and J.L. Meunier, Nucl. Phys. B207, 251 (1982).

[12] D. Appel, P. Mackenzie, and G. Sterman, Nucl. Phys. B309, 259 (1988).

[13] H. Contopanagos and G. Sterman, Nucl. Phys. B400, 211 (1993); B419, 77 (1994).

[14] L. Alvero and H. Contopanagos, Nucl. Phys. B436, 184 (1995); B456, 497 (1995).

[15] E. Laenen, J. Smith, and W.L. van Neerven, Nucl. Phys. B369, 543 (1992); Phys. Lett. B 321, 254 (1994).

[16] E.L. Berger and H. Contopanagos, Phys. Lett. B 361, 115 (1995); Phys. Rev. D 54, 3085 (1996); ANL-HEP-CP-96-51, hep-ph/9606421, 1996.

[17] S. Catani, M.L. Mangano, P. Nason, and L. Trentadue, Nucl. Phys. B478, 273 (1996); Phys. Lett. B 378, 329 (1996).

[18] N. Kidonakis and J. Smith, Phys. Rev. D 51, 6092 (1995); Mod. Phys. Lett. A 11, 587 (1996); ITP-SB-95-16, hep-ph/9506253, 1995.

[19] N. Kidonakis, J. Smith, and R. Vogt, hep-ph/9608343, 1996.

[20] D.B. DeLaney, S. Jadach, Ch. Shio, G. Siopsis, and B.F.L. Ward, Tennessee preprint UTHEP-95-0102, 1995.

[21] J.C. Collins, D.E. Soper, and G. Sterman, Nucl. Phys. B250, 199 (1985);
C.T.H. Davies and W.J. Stirling, Nucl. Phys. B244, 337 (1984);
C.T.H. Davies, B.R. Webber, and W.J. Stirling, Nucl. Phys. B256, 413 (1985);
P.B. Arnold and R.P. Kauffman, Nucl. Phys. B349, 381 (1991);
G. Ladinsky and C.-P. Yuan, Phys. Rev. D 50, 4239 (1994);
Yu.L. Dokshitzer, D.I. D'Yakonov, and S.I. Troyan, Phys. Lett. 79B, 269 (1978);
G. Parisi and R. Petronzio, Nucl. Phys. B154, 427 (1979).

[22] J.C. Collins and D.E. Soper, Nucl. Phys. B193, 381 (1981); B197, 446 (1982);
C.L. Basham, L.S. Brown, S.D. Ellis, and S.T. Love, Phys. Rev. D 19, 2018 (1979);
L.S. Brown and S.D. Ellis, Phys. Rev. D 24, 2383 (1981).

[23] A.M. Polyakov, Nucl. Phys. B164, 171 (1979);
I.Ya. Aref'eva, Phys. Lett. 93B, 347 (1980);
V.S. Dotsenko and S.N. Vergeles, Nucl. Phys. B169, 527 (1980);
R.A. Brandt, F. Neri, and M.-a. Sato, Phys. Rev. D 24, 879 (1981).

[24] J. Botts and G. Sterman, Nucl. Phys. B325, 62 (1989).

[25] G. P. Korchemsky, Phys. Lett. B 325, 459 (1994).

[26] I. A. Korchemskaya and G. P. Korchemsky, Nucl. Phys. B437, 127 (1995).

[27] G.P. Korchemsky and G. Sterman, Nucl. Phys. B437, 415 (1995).

[28] N. Kidonakis and G. Sterman, Phys. Lett. B 387, 867 (1996); in *Les Rencontres de Physique de la Vallée d'Aoste, Results and Perspectives in the Standard Model*, La Thuile, Aosta Valley, March 4-9, 1996, ed. M. Greco (Laboratori Nazionali de Frascati SIS-Ufficio Pubblicazioni, Frascati, Italy, 1996), p. 333, hep-ph/9607222.

[29] N. Kidonakis, SUNY at Stony Brook Ph.D. Thesis, hep-ph/9606474, UMI-96-36628-mc (microfiche), 1996.

[30] S. Dawson, Nucl. Phys. B359, 283 (1991);
A. Djouadi, M. Spira, and P.M. Zerwas, Phys. Lett. B 264, 440 (1991);
M. Spira, A. Djouadi, D. Graudenz, and P.M. Zerwas, Nucl. Phys. B453, 17 (1995);
R.P. Kauffman, Phys. Rev. D 45, 1512 (1992);
S. Dawson and R.P. Kauffman, Phys. Rev. D 49, 2298 (1994);
C.-P. Yuan, Phys. Lett. B 283, 395 (1992);
M. Krämer, E. Laenen, and M. Spira, CERN-TH-96-231, hep-ph/9611272, 1996.

[31] E. Laenen and G. Sterman, in *Proceedings of The Fermilab Meeting, DPF 92*, 7th meeting of the American Physical Society Division of Particles and Fields (Batavia, IL, 1992), ed. C. H. Albright *et al* (World Scientific, Singapore, 1993), p. 987.

[32] G.P. Korchemsky and G. Marchesini, Nucl. Phys. B406, 225 (1993);
G.P. Korchemsky and G. Sterman, Phys. Lett. B 340, 96 (1994);
R. Akhoury and I.Z. Rothstein, Phys. Rev. D 54, 2349 (1996).

[33] J. C. Collins, D. E. Soper, and G. Sterman, in *Perturbative Quantum Chromodynamics*, ed. A. H. Mueller (World Scientific, Singapore, 1989), p. 1.

[34] J. G. M. Gatheral, Phys. Lett. B 133, 90 (1983);
J. Frenkel and J.C. Taylor, Nucl. Phys. B246, 231 (1984);
G.P. Korchemsky and A.V. Radyushkin, Phys. Lett. B 171, 459 (1986);
Nucl. Phys. B283, 342 (1987);
J. C. Collins, in *Perturbative Quantum Chromodynamics*, ed. A. H. Mueller (World Scientific, Singapore, 1989), p. 573.

[35] L. N. Lipatov, Nucl. Phys. B309, 379 (1988).

[36] R. Meng, G.A. Schuler, J. Smith, and W. L. van Neerven, Nucl. Phys. B339, 325 (1990).

[37] M. Sotiropoulos and G. Sterman, Nucl. Phys. B419, 59 (1994).

[38] B.R. Webber, Phys. Lett. B 339, 148 (1994);
A.V. Manohar and M.B. Wise, Phys. Lett. B 344, 407 (1995);
Yu.L. Dokshitzer and B.R. Webber, Phys. Lett. B 352, 451 (1995);
R. Akhoury and V.I. Zakharov, Phys. Lett. B 357, 646 (1995);
P. Nason and M.H. Seymour, Nucl. Phys. B454, 291 (1995);
M. Beneke and V.M. Braun, Nucl. Phys. B454, 253 (1995);
M. Beneke, hep-ph/9609215, 1996.